



Development of a video-microscopic method to compare the effect of a precipitation inhibitor

Christfort, J.F.; Plum, J.; Madsen, C.M.; Nielsen, Line Hagner; Müllertz, A.; Rades, T.

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Christfort, J. F., Plum, J., Madsen, C. M., Nielsen, L. H., Müllertz, A., & Rades, T. (2016). *Development of a video-microscopic method to compare the effect of a precipitation inhibitor*. Poster session presented at 2016 AAPS Annual Meeting and Exposition, Denver, CO, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


PURPOSE

The aim of this study was to develop a video-microscopic method to evaluate the effect of a precipitation inhibitor (PI) on supersaturated solutions of the poorly soluble drug tadalafil using a novel small scale setup.

CONCLUSION

- Tadalafil shows a prolonged induction time and a reduced growth rate in presence of HPMC.
- To significantly prolong the induction time and decrease particle growth, 0.01 % w/v HPMC is needed.
- This is a promising tool for evaluating the effect of PI's on induction time and crystallization rate of supersaturated systems of poorly soluble drugs.

METHOD



ACKNOWLEDGEMENTS


Philips BioCell A/S is acknowledged for support and access to the equipment.

This study was conducted as part of the Oral Biopharmaceutics Tools (ORBITO) project (<http://www.orbitoproject.eu>), funded by the Innovative Medicines Initiative (IMI) Joint Undertaking under Grant Agreement No. 115369.




RESULTS

Proof of Concept using the commercial software




Top: Induction time for tadalafil in presence of HPMC
Bottom: The area of one well defined particle per well as a function of time, with different HPMC conc., mean \pm SD, n=3-24

Segmentation



- 0.0 % (w/v) HPMC
- 0.001 % (w/v) HPMC
- ▲ 0.01 % (w/v) HPMC
- ▼ 0.1 % (w/v) HPMC
- ◆ 0.5 % (w/v) HPMC

Improved Image analysis



Top: Induction time for tadalafil in presence of HPMC
Bottom: The total particle area as a function of time, with different HPMC conc., mean \pm SD, n=24